

FUNGI NONNULLI EXTRA-ITALICI NOVI

EX HERBARIIS

C. C. GILLET, P. MORTHIER, ET G. WINTER.

AUCTORE

P. A. SACCARDO

1. **Pseudopeziza Morthieri** Sacc. Maculis foliorum are-
scendo ochraceo-fulvis, amplis; cupulis hypophyllis hinc
inde gregariis, plano-scutellaribus, adnato - sessilibus,
minutissimis, vix $\frac{1}{10}$ mill. diam., glabris, madore disco
apertis, flavo-fulvis, pellucidis; contextu excipuli paren-
chymatico, margine subprosenchymatico; ascis cylindra-
ceo-clavulatis, 45 \times 7, subsessilibus, paraphysibus fili-
formibus apice *aduncis*, obvallatis, octosporis; sporidiis
distichis, cylindraceo-clavatis, 7-10 \times 2-2 $\frac{1}{2}$, quandoque
curvulis, hyalinis.

Hab. in pagina inferiore foliorum Rubi glandu-
losi in alpibus Helveticis, unde misit Junio 1878 cla-
riss. prof. Doct. P. MORTHIER.

Obs. A *Peziza misella* Dmz., *P. dumorum* Dmz.,
P. Edema Dmz. nec non a *Pseudopeziza exigua* Niessl
(omnibus folia Ruborum habitantibus) probe diversa.

2. **Pleospora Gilletiana** Sacc. Peritheciis gregariis, epi-
dermide tumidula velatis, e globoso depressis, $\frac{1}{2}$ mill.
diam., nigris, vix papillatis, basi hyphis filiformibus co-
piosis, fuligineis cinctis, ceterum glabris; contextu laxe
parenchymatico, fuligineo; ascis cylindraceis 130 \times 13-
15, subsessilibus, apice rotundatis ibique lumine trun-
cato integro, paraphysibus (ubi adsunt) brevissimis ob-
vallatis, octosporis; sporidiis oblique monostichis ob-
longo-ovoideis, medio plerumque leniter constrictis,
utrinque obtusiusculis, 25-28 \times 12-13, initio hyalinis

1-septatis, dein 7-septatis denseque muriformibus, cribrose guttulatis, flavis, tandem melleis.

Hab. in ramulis *Genistæ* prope *Alençon* in Gallia 1878 (C. C. GILLET).

Obs. A *Pleospora Cytisi* Fckl., cui affinis, differt ascis multo angustioribus, sporidiis brevioribus, peritheciis basi vestitis &c.

Lophiotrema Winteri Sacc. Peritheciis sparsis gregariisve, cortice nidulantibus lignoque adnatis, globulosis, $\frac{1}{2}$ mill. d., ostiole emergente compresso sed angusto et acutiusculo; ascis clavatis, v. clavato-cylindraceis, 100–110 \times 15–17, apice rotundatis, breve stipitatis, paraphysibus copiosis filiformibus obvallatis, octosporis; sporidiis monostichis vel p. p. distichis, fusoideis, curvulis, 28–34 \times 7 $\frac{1}{2}$ –8 $\frac{1}{2}$, obtusiusculis, 5-septatis, ad septa leniter constrictis, typice hyalinis 6-guttulatis, utrinque appendiculâ longiusculâ crassâ hyalinâ, mox deciduâ auctis.

Hab. in ramulis emortuis *Helianthemi* prope *Zurich* in *Helvetia*, aut. 1878 (Prof. G. WINTER).

Obs. Peraffine *L. præmorso* (Lasch.) Fckl., Sacc. F. it. tab. 253, a quo peritheciis minoribus, ostiolis acutiusculis, sporidiis crassius appendiculatis recedit et *L. sexnucleato* Cke a quo sporidiis appendiculatis, ostiole angustiore &c. differt.

3. **Phoma aculeorum** Sacc. Peritheciis gregariis, punctiformibus, $\frac{1}{10}$ – $\frac{1}{8}$ mill. d., conoideis, nigris, initio epidermide velatis; spermatiis cylindraceo-oblongis, saepe curvulis, 3–4 \times 1, obsolete 2-guttulatis, hyalinis, basidiis fusoideis crassiusculis suffultis.

Hab. in aculeis languidis *Rosarum* prope *Alençon* in Gallia, 1878 (C. C. GILLET).

4. **Phoma enteroleuca** Sacc. Peritheciis hinc inde acervulatis, peridermio tectis, dein erumpentibus, globulosis, $\frac{1}{4}$ mill. diam. vix papillatis, diu nucleo albo sarcis; spermatiis ovoideo-oblongis, 4 \times 1 $\frac{1}{2}$, eguttulatis, hyalinis (basidiis nullis visis).

Hab. in ramulis *Pyri* prope *Alençon* in Gallia, 1878 (C. C. GILLET).

5. **Phoma populea** Sacc. Botryosphaeroides; peritheciis

hinc inde aggregatis, innato-erumpentibus, globulosis, minutis, nigris, vix papillatis; spermatiis obovoideis, 10 \times 7, apice rotundatis, nubilosus, hyalinis (basidiis nullis visis).

Hab. in ramis corticatis Populi fastigiatæ prope Alençon in Gallia, 1878 (C. C. GILLET).

6. **Phoma xanthina** Sacc. Peritheciis gregariis, epidermide tumidula velatis, globoso-depressis, $1\frac{1}{5}-1\frac{1}{3}$ mill. diam., ostiolo exiguo erumpente, contextu laxe parenchymatico fuligineo; spermatiis cylindraceis v. cylindraceo-oblongis, utrinque obtusis, curvulis, 15 \times 4-6, 2-3-guttulatis, singulis subhyalinis, coacervatis luteolis, basidiis obsoletis.

Hab. in caule putrescente Delphinii prope Alençon in Gallia, 1878 (C. C. GILLET).

7. **Phoma Spartii** Sacc. Peritheciis gregariis, cortice velatis, dein, ipso secesso, superficialibus, basi ligno insculptis, oblongis, quandoque subhysteriformibus, $\frac{1}{3}$ mill. lgs., $\frac{1}{5}$ crs., nigris, ostiolo saepe rimose dehiscente; spermatiis fusoideis, 10-11 \times 2- $2\frac{1}{2}$, 2-guttulatis, hyalinis, basidiis filiformibus, curvis, 20-22 \times $1\frac{1}{2}$ -2, suffultis.

Hab. in trunco Spartii prope Alençon, in Gallia 1878 (C. C. GILLET).

Obs. Verisimiliter status spermogonicus *Diaporthes* cuiusdam.

8. **Vermicularia herbarum** West., Kx Fl. Flandr. I. 405 (ut videtur).

Hab. in caule putrescente Solani tuberosi prope Alençon in Gallia, 1878 (C. C. GILLET).

Obs. Perithecia laxiuscule gregaria, superficia, globoso-depressa, $\frac{1}{2}$ mill. diam., nigra, setulis crebris vestita; spermatia cylindracea, subrecta, utrinque obtuse rotundata, 20-22 \times 3-4, granulosa, hyalina. A *Verm. Dematio* Fr., cui habitu affinis, differt spermatiis rectis, obtusisque, nec falcatis. Longius distat a *Vermic. atramentaria* B. & Br. (? == *V. maculanti* [Link] Dmz.) ob perithecia majora & laxe sparsa.

9. **Septoria Gilletiana** Sacc. Maculis nullis; peritheciis

(spuriis) hypophyllis, hinc inde gregariis 100–110 micr. diam. epidermide tumidula tectis; spermatiis e strato proligero crassiusculo fasciculatim oriundis, filiformibus curvulis v. tortuosis, 30–45 × 1 $\frac{3}{4}$ –2, utrinque obtusiusculis, 3-septatis, hyalinis.

Hab. in foliis nondum emortuis *Castaneæ* vescæ prope *Alençon* in Gallia, 1878 (C. C. GILLET).

Obs. A *Sept. castanicola* Dmz., Sacc. *Michelia* II. 175. spermatiis dimidio et ultra angustioribus &c. recedit.

10. **Phyllosticta Platanoidis** Sacc. Maculis subnullis, peritheciis minutissimis, hypophyllis, hinc inde maculiformiter aggregatis, vix 70–80 micr. diam., poro pertusis; spermatiis oblongis, perexilibus, 3 × $\frac{1}{2}$, hyalinis.

Hab. in foliis *Aceris Platanoidis* prope *Alençon* in Gallia, 1878 (C. C. GILLET).

11. **Glæosporium phæosorum** Sacc. Acervulis ramulicolis, gregariis epidermide velatis, globosis v. oblongis, $\frac{1}{4}$ mill. diam., nigricantibus, tandem irregulariter erumpentibus; conidiis fusoideis majusculis, 25–28 × 8, raro oblongis, chlorino-hyalinis granulosis, basidiis filiformibus, brevibus, quandoque fuscatis fultis.

Hab. in sarmentis corticatis *Ruborum* prope *Alençon* in Gallia, 1878 (C. C. GILLET).

Obs. Äquo fere jure ad gen. *Myxosporium* duci potest.

12. **Stemphylium ramulosum** Sacc. Effusum, velutinum, atrum, hyphis ascendentibus, filiformibus, elatis, $\frac{1}{2}$ mill. altis, 10–13 micr. crassis, deorsum leniter incrassatis, apice simpliciter v. repetitive breve-ramulosis, ubique crebre articulatis, intense fuligineis, articulis 2-guttatis; conidiis ex apice hypharum oriundis, oblongis v. obpyriformibus, 35–50 × 18, 5–7septato-muriformibus, cibrose guttulatis, fuligineis,

Hab. in caule putrescente *Apri Petroselini* prope *Alençon* in Gallia (C. C. GILLET).

Obs. Genus *Mystrosporium* a *Stemphylio* antiquiore essentialiter non differt. Species haec a *Mystr. Spraguei* B. & Curt. et *Mystr. macropode* Cda. hyphis apice ramosis statim dignoscitur.